

TO ALL TO VILONITHESE PRESENTS SHAML COME; Yornell University Agricultural Experiment Station Wilherens, There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, upon due examination made, the said applicant(s) is (are) adjudged TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF Seventeen YEARS FROM THE DATE OF THIS GRANT, SUBJECT O THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC ED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-UDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS

BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

ALFALFA

'Multileas'

In Bestimony Watercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 16th day of March / in the year of our Lord one thousand nine hundred and seventy-eight

NSTRUCTIONS: See Reverse.	2. KIND NAME	<u> </u>		IAL USE ONLY
VARIETY NAME OR TEMPORARY DESIGNATION			PV NUMBER	
'Multileaf'	Alfalfa		7310	
	4. FAMILY NAME (Bota	micul)	FILING DATE	TIME A.M. A.M.
3. GENUS AND SPECIES NAME		•	6.18.73	11:00-
Medicago sativa	Leguminosae		FEE RECEIVED	BALANCE DUE
	5. DATE OF DETERM	NATION	3.50.00	\$6+18/13
	11 October 19	72	\$ 050.00	- Chilaran
			1: 350.00	B. TELEPHONE AREA
6. NAME OF APPLICANT(S)	7. ADDRESS (Street an	d No. or R.F.D. No.	, City, State, and ZIP	CODE AND NUMBER
Cornell University	Cornell Unive	ersity		(07 05/ 5/00
Agricultural Experiment	Ithaca, New Y	ork 14853		607-256-5420
Station Agricultural Experiment				
		•		
	ERSON, FORM OF	10. STATE OF INC	ORPORATION	11. DATE OF INCOR-
9. IF THE NAMED APPLICANT IS NOT A PE ORGANIZATION: (Compration, partnership	, association, etc.)			
		New York		1888
State Agricultural Experim 12. Name and mailing address of appli	icant representative(s), if any, to serve	e in this application	and receive all papers
Mame and marring address of -bb.	•	•		•
R. P. Murphy	·	•		
Department of Plant Breedi	ing and Biometry		•	
Cornell University			607_256_3101	
Ithaca, New York 14853	•	Telephone	607-256-3101	
		·	<u>,,</u>	
13. CHECK BOX BELOW FOR EACH ATTA	CHMENT SUBMITTED:	Varior / 5 5	tion 52 of the Plant	Variety Protection Act
13. CHECK BOX BELOW FOR EACH ATTAG	reeding History of the	variety (see sec		•
[X] 138. Exhibit B, Botanical De	scription of the varie	-7		
*				
X 13c. Exhibit C, Objective De	acription of the fact	- To the second		
[X] 130. Exhibit D, Data Indicati	ive of Noveltv			
•	•			
X 136. Exhibit E, Statement of	the Basis of Applican	nt's Ownership		
14A. Does the applicant(s) specify t	har send of this varie	ty be sold by var	iety name only as a	class of certified seed
(See Section 83(a), (If "Yes,"	answer 14B and 14C	below.)	XYES D	10
(See Section 83(a), (1), 183,	answer 140 and 110	14C. If "Yes,"	to 14B, how many g	generations of production
148. Does the applicant(s) specify t	tions?	beyond br	eeder seed?	
limited as to number of general	IXIYES I INC	У ГОПИО	ATION REGISTE	
The applicant declares that a viable	L. Chapia se	ad of this variety	will be deposited ut	oon request before issu
The applicant declares that a viable ance of a certificate and will be re	re sample of basic sec	v in accordance:	with such regulations	s as may be applicable
	seleesehed beriodicali	V III accordance	W **** C ***	·
The undersigned applicant(s) of	this sexually-reproduc	rea novet plant v	on under the transei	ons of Section 42 of th
uniform, and stable as required in	n Section 41 and is e	ntitiea to protecti	ton unuer the propest	
Disea Marioty Protection Act				
Applicant is informed that false	representation herein	can jeopardize p	rotection and result	m penaintes.
••••		\sim	11 10	<u>, </u>
	,	\Box :	+ husto	4
Mar h V3 197	<u> </u>	—— —	(SIGNATURE OF AP	WICANT)

7 (SIGNATURE OF APPLICANT) (DATE)

- - - - -

1

13A. Exhibit A

Origin and Breeding History of the Variety

- 1. 'Multileaf' originated from research in the Department of Plant Breeding and Biometry, Cornell University Agricultural Experiment Station, New York State College of Agriculture, Cornell University, Ithaca, New York. The original plants were selected for extra leaflets per leaf from the breeding nurseries. The first population of plants was produced from eleven plants in the nurseries each of which exhibited a low level of expression of the multifoliolate character. The origin of these plants was: five from the breeding population that produced 'Saranac', three from the breeding population that produced 'Iroquois', two from the cultivar 'Narragansett', and one of unknown origin from an old field in New York. Five cycles of recurrent selection for extra leaflets per leaf were made since 1959.
- 2. Selection has also been based on vigor, winter survival and resistance to the bacterial wilt disease. All selections have been made in field nurseries.
- 3. One hundred parent clones were interpollinated to produce breeder seed of this cultivar.
- 4. The within cultivar uniformity is similar to that for others developed at this experiment station, i.e. 'Saranac' and 'Iroquois', and 'Vernal' and 'Lahontan' and meets the uniformity requirements of alfalfa cultivars. See Exhibit C. The most distinguishing characteristic is the multifoliolate character; we expect less than one percent of the population to be trifoliolate. Not all leaves on a multifoliolate plant exhibit the character; individual plants may vary from 50 to 85 percent of their leaves having more than three leaflets. The percentage of multifoliolate plants in the cultivar has remained stable through two generations beyond the Breeder seed generation.

Botanical Description of the Variety

'Multileaf' is typical of <u>Medicago sativa</u> except for the expression of the multifoliolate character (more than three leaflets per leaf). The seed pods are coiled. The flower color is characterized as follows: 46% light through dark purple, 40% light through dark reddish purple, 12% purple to blue as flowers age, and 2% purple to some variegation as flowers age.

Objective Description of the Variety

'Multileaf' alfalfa is a winter-hardy cultivar under the climatic conditions of Central New York. It is resistant to the bacterial wilt disease. Growth type, regrowth characteristics and fall dormancy reaction are similar to those of the cultivar 'Saranac'. Performance information indicates total-season dry matter production of 'Multileaf' to be 90-95% of that for the 'Saranac' and 'Iroquois' cultivars. Determinations of leaf percentage of total dry matter have indicated 'Multileaf' exceeds 'Saranac' by 2 to 3% for forage managed and harvested under the same conditions.

13D. Exhibit D

Data Indicative of Novelty ('Multileaf' 73100)

Novelty is based on the characteristic of multifoliolate leaves (more than three leaflets per leaf). The most similar cultivars are Iroquois and Saranac.

Cultivar	Percent of plants with multifoliolate leaves	Percent of leaves on individual plants with more than three leaflets			
'Multileaf'	99	55 to 85			
'Iroquois'	>1	>1			
'Saranac'	>1	>1			

Multifoliolate leaves have 5 leaflets most frequently. Expression of the multifoliolate leaf character appears to be somewhat greater on forage produced during the cooler and shorter days of early spring and late fall than in mid-summer.

'Multileaf' may be distinguished from all cultivars known to us very clearly because of its multifoliolate leaf character. See Exhibit C. 'Multileaf' is the only cultivar known to us which exhibits the multifoliolate leaf character.

Data from a bacterial wilt test on 10 varieties tested with two sources of inoculum (Minnesota and New York). There were no variety x source of inoculum interactions. Conducted in the field in 1975 at Ithaca, N. Y. by R. P. Murphy.

Variety	Wilt Score	(Reaction and survival)
Agate	. 0.77	
Iroquois	1.14	
Saranac AR	1.17	
Vernal	1.34	
Saranac	1.45	
Apollo	1.45	
Honeoye	1.66	
Ranger	1.81	
WL-305	1.89	
DuPuits	3.35	

All of these varieties are described as resistant except DuPuits.

Data from anthracnose test on varieties included in the 1974 Alfalfa Variety Test sponsored by NE-74 Regional Research Project (Courtesy U.S. Regional Pasture Research Laboratory).

<u>Variety</u>	Anthracnose Score	(0=immune to 5=dead)
Saranac AR	2.67	
Arc	2.68	
Ramsey	3.64	
Agate	3.81	
TitAn	4.18	
Aztec	4.40	
530	4.59	
520	4.62	
Honeoye	4.64	
Victoria	4.68	
Multileaf	4.78	
Iroquois	4.81	
Saranac	4.84	
Kodiak	4.88	
Bonus	4.98	

This test was conducted in a plant growth chamber and at a very high level of infection. Data recorded on 4/7/75.

Observations from variety yield trials on the multifoliolate character:

None of the following varieties show a degree of multifoliolate plants as does Multileaf ($\pm 99\%$ of plants with some degree of this character). A trace of such plants ($\pm 1\%$) may be found in Narragansett.

Agate Phytor Alfa Ramsey Americana Team Anchor Thor Apollo Titan Vernal ArcAztec Victor Vista Bonus Cardinal Weevlchek CW-5 WL-218 Dawson WL-219 WL-305 DuPuits Gemini WL-307 Glacier WL-309 WL - 311Kanza Klondike WL-318 Kodiak 131 167 Nugget Narr**A**gansett 520 Olympian 521 Pacer 530

Observations from variety yield trials on flower color:

None of the following varieties show a significant degree of variegated or yellow flowered plants as does Honeoye ($\pm 85\%$ of plants with variegated or yellow flowers).

Alfa	Phytor
Anchor	Team
Apollo	Thor
Arc	Victor
Aztec	Vista
Bonus	WL-219
Cardinal	WL-305
CW-5	WL-307
Dawson	WL-309
DuPuits	WL-311
Kanza	WL-318
Klondike	131
Kodiak	520
Nugget	521
Olympian	530

13E. Exhibit E

Statement of Applicant's Ownership

The Cornell University Agricultural Experiment Station is the owner of 'Multileaf'.

FORM GR 470-32 (3/75)

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION HYATTSVILLE, MARYLAND 20782

EXHIBIT C

OBJECTIVE DESCRIPTION OF VARIETY Alfalfa (Medicago sativa L. complex)

	VARIETY NAME OR TEMPORARY
NAME OF APPLICANT(S)	DESIGNATION Multileaf
NAME OF APPLICANT(S) Cornell University Agricultural Experiment Station Cornell University Agricultural Experiment Station	FOR OFFICIAL USE ONLY
ADDRESS (Street and No., of the second TV	PVPO NUMBER
Department of Plant Breeding and Biometry New York State College of Agriculture and Life Sciences New York 14853	73100
New York State College of Agricultury Vork 14853	<u> </u>
Cornell University, Ithaca, New 1 hands of this variety in the boxes	s below.
and the number that describes the varietal character and the constant of the c	or less.
Di GOEO ID FIFST DOX (C.E. 10 10 10 10 10 10 10 10 10 10 10 10 10	
NOTE: For single plant data a minimum of 100 plants is suggested	INDICATE AREA WHERE TEST WAS
1. PRIMARY AREA OF ADAPTATION	CONDUCTED. FURTHER EXPLANATION CAN GO IN COMMENTS AT THE END
	OF THE FORM.
1 = NORTHWEST 2 = NORTHCENTRAL 3 = NORTHEAST	3 AREA TESTED
S = SOUTHERN PLAINS	L31 AMEA 13-1
3 4 = SOUTHEAST 5 = SOUTHWEST 6 = SOUTHWEST	
2. WINTER HARDINESS	\
1 = NON-HARDY (Mesa Sirsa) 3 = INTERMEDIATE NON-HARDY 1 = NON-HARDY (Mesa Sirsa) 7 = HARDY (Vernal)	3 AREA TESTED
5 = MODERATELY HARDY (Saranac) 9 = EXTREMELY HARDY (Norseman)	
g-EATHERS .	
1 = ANTICIPATED 2 = MEASURED	l .
2 SOURCE OF INFORMATION: 1 = ANTICIPATED 2 = MEASURED	
3. FALL GROWTH HABIT	
a - SEMIERECT (DuPuits)	3 AREA TESTED
1 = ERECT (Mesa Sirsa) 3 = SEMIERECT (Datation) 5 = INTERMEDIATE (Saranac) 7 = SEMIDECUMENT (Vernal)	3 AREA TESTED
9 = DECUMBENT (Norsement)	
4. RECOVERY AFTER FIRST SPRING CUTTING	
5 = INTERMEDIA	TE C
1 = VERY FAST (Mesa Sirsa) 3 - 745 (Old (Norseman)	3 AREA TESTED
3 7 = SLOW (Vernal)	
5. FLOWERING DATE (FIRST SPRING GROWTH)	·
	\ 1
1 = MESA SIRSA 2 = LAHONTA 3 = SARANAC 4 = VERNAL	AN 3 AREA TESTED
10131	
0 0 DAYS LATER THAN 3 5 = NORSEMAN	
 * L - 	
6. CROWN TYPE	\
1 = SPREADING ROOTS 3 = SPREADING RHIZOMES (Teton)	3 AREA TESTED
7 = INTERMEDIATE (Saranac)	
9 = NARROW (Mesa Sirsa)	
7. PLANT COLOR	_
3 = DARK GREEN (Weevichek) 5 = GREEN (Vernal)	3 AREA TESTED
4 7 = LIGHT GREEN (Ranger)	
8. HAIRINESS	
Г	% PLANTS WITH PUBESCENT
% PLANTS WITH PUBESCENT STEMS	PODS
<u></u>	prop 1
9, POD SHAPE	5
	OUS 0 0 0 % PLANTS WITH SICK
0 9 8 % PLANTS WITH TIGHT COILS 0 0 2 % PLANTS WITH LOOSE C	PODS (Less than 1 coil)
0 9 8 % PLANTS WITH TIGHT COILS 10111	

FORM GR-470-32 (2	2/75)							Ţ	310	Ö PA	GE 2 OF 4
10. GIVE ITEM LEI	NGTH FREQUENC	Y DISTRIE	UTION F	OR SUBM	ITTED A	VD 1 TO 5	STANDA	RD VARI	ETIES 1	!	
•		1 1			UENCY D	ISTRIBUT	ION 2/	· 			
VARIETY NAME	O – 5 6 – 10 mm, mm, % %	11 — 15 mm. %	16 – 20 mm. %	21 — 30 mm. %	31 – 40 mm, %	41 — 50 mm. %	51 – 60 mm. %	61 — 70 mm. <u>%</u>	71 – 80 mm. =- %	81 + mm, %	AVERAGE STEM LENGTH
Multileaf	46.0		37.2	12.9	2,6	0.5	0.5	0.0	0.0	0.0	13.40
Lahontan	19.8	ļ <u>.</u>	48.9	20.4	7.7	2.3	0.0	0.2	0.2	0.0	18.10
Saranac	69.7		24.8	4.8	0.5	0.0	0.0	0.0	0.0	0.0	10.00
Iroquois	71.7		23.8	3.8	0.2	0.0	0.2	0.0	0.0	0.0	9.80
Vernal	82.5		14.6	2.5	0.0	0.2	0.0	0.0	0.0	0.0	8.60
11. FLOWER COLO	R 3/ (DETERMIN	E COLOR (ON FRES	HLY OPE	NED FLO	NERS)				<u></u>	
	PURPLE 0 0	2 % VA		1	0 0	% YELLOW	00	0 %	CREAM	0 0	0 % WHITE
12. DISEASE, INSE		-				•			-		
DISEASE	CULTIVA		% RES	STANT		VERITY	ASI LSD .6	1			ATION _4/
	Multileaf		23		2.3		.42				Minnesot
	(SUBMITTED)	 			<u> </u>	4		1974, Rosemont, Minn. Field			
BACTERIAL WILT	(BES, CK.) VERN	35	35.1		.4		Fi				
	(SUS, CK.) NARR	AG ANSET	- 00	.0	4.3	10					
	Multileaf			4.7		.48	U.	U.S. Regional Pasture			
	(SOBMITTED)							Research Laboratory, 1975, State College, Pa. Greenhouse			
ANTHRACNOSE	(RES, CK.) ARC			2.6	2.68						
	(SUS. CK.) SARA	VAC			4.8	4					
	(SUBMITTED)				. _						
COMMON LEAF SPOT	(RES. CK.) RAMS	ĒΥ	<u>L.</u> .								
SFU1	(SUS. CK.) RANG	ER					<u> </u>				
	(SUBMITTED)		<u> </u>		_						
DOWNY MILDEW	(RES, CK.) SARANAC		<u> </u>								. .
	(SUS. CK.) KANZ	Α									
	(SUBMITTED)	-									
PHYTOPHTHORA ROOT ROT	(RES. CK.) AGAT	E									
	(SUS. CK.) SARAI	NAC				· <u>.</u>	ļ			 	
	(SUBMITTED)		 								
OTHER	(RES. CK.)		<u> </u>								
	I		ſ				1				

(SUS. CK,)

^{1/} Preferred standards: Saranac, Vernal, Norseman, Lahontan, Mesa Sirsa. Twelve hours light at 25° C with 20,000 lux of cool white florescent; 2,000 lux of incandescent filament light and twelve hours darkness at 5% C.
2/ From cotyledonary node to tip of stem 20 days after planting.
3/ For further clarification consult USDA Agricultural Handbook No. 424.
4/ Give: The institution in charge of test, (2) year, and (3) location of test. Describe test procedure if it differs from procedure suggested in ARS-NC-19, September 1974.

12. DISEASE, INS	ECT, AND NEMATODE RESI	STANCE: (Enter re		nd check cult	ivars. Circle check cultivars used.)
DISEASE	CULTIVAR	% RESISTANT PLANTS	AVG. SEVERITY INDEX (ASI)	ASI LSD ,05	TEST, YEAR & LOCATION 4/
	(SUBMITTED)				
				1	
OTHER	(RES, CK.)			1	
	(SUS. CK.)				· · · · · · · · · · · · · · · · · · ·
	(SUBMITTED)				
OTHER	(RES. CK.)				
	(SUS. CK.)	,			
INSECT	CULTIVAR	% SEEDLING SURVIVAL	AVG' SEVERITY INDEX (ASI)	ASI LSD .05	TEST, YEAR & LOCATION 4/
 :	(01)-117750)				
	(SUBMITTED)				
PEA APHID	(RES. CK.) KANZA				
	(SUS. CK.) RANGER	-			
	(SUBMITTED)			1	
SPOTTED	(RES. CK.) KANZA]	
ALFALFA APHID	(HED, DKJ) KANEA			[
	(SUS, CK.) RANGER	i I	AVG. SEVERITY	ASI	
INSECT	CULTIVAR	% DEFOLIATION	INDEX (ASI)	LSD .05	TEST, YEAR & LOCATION 4/
	(SUBMITTED)		·		
ALFALFA WEEVIL	(RES. CK!) ARK				
	(SUS. CK.) VERNAL				
INSECT	CULTIVAR	% RESISTANT PLANTS	EMERGED ADULTS PER PLANT	EMERGED LSD .05	TEST, YEAR & LOCATION 4/
	(SUBMITTED)		,		
	12.00		.,	1 1	
ALFALFA SEED CHALCID	(RES. CK.) LAHONTAN				
	(SUS. CK.) SONORA	a, pp	AMO SELECTION		
INSECT	CULTIVAR	% RESISTANT PLANTS	AVG, SEVERITY INDEX (ASI)	ASI LSD .05	TEST, YEAR & LOCATION 4/
	(SUBMITTED)]	
POTATO LEAF-	(RES. CK.)				
HOPPER				1 1	
	(SUS. CK.)			 	
i	(SUBMITTED)				
OTHER	(RES. CK.)		<u> </u>] [
'				I	

⁽SUS. CK.)

4/ Give: The institution in charge of test, (2) year, and (3) location of test. Describe test procedure if it differs from procedure suggested in ARS-NC-19, September 1974. F

12. DISEASE, INS	ECT, AND	NEMATODE RESIS	STANCE: (Enter resis	tance of submitted		tivars. Circle check cultivars used.)	
INSECT		CULTIVAR	% RESISTANT PLANTS	AVG, SEVERITY INDEX (ASI)	ASI LSD ,05	TEST, YEAR & LOCATION 4/	
OTHER	(SUBMI	TTED)					
	(RES. C	K.)					
·	(sus, c	K.)			ASI		
NEMATODE		CULTIVAR	% RESISTANT PLANTS	INDEX (ASI)	ASI LSD .05	TEST, YEAR & LOCATION 4/	
	(SUBM	ITTED)					
STEM NEMATODE	(RES. C	K.) LAHONTAN			<u> </u>		
	(SUS. C	K.) RANGER					
	(SUBM	ITTED)					
NORTHERN ROOT KNOT	(AES. C	CK.) NEV. SYN. XX	-		i		
NEMATODE	(SUS. C	CK.) LAHONTAN					
	(SUBM	ITTED)					
SOUTHERN ROOT KNOT	(RES.	CK.) MOAPA 69					
NEMATODE	(SUS. (CK.) LAHONTAN					
	(SUBM	ITTED)					
OTHER	(RES.	CK.)					
	(SUS.						
13. INDICATE A	VARIET	Y THAT MOST CLOS	SELY RESEMBLES T	HE VARIETY SUB	MITTED FOR	THE FOLLOWING CHARACTERS:	
CHARACTER VARIETY			CHARAC	TER	VARIETY		
AREA OF ADAP	AREA OF ADAPTATION Saranac			PLANT HEIGH	PLANT HEIGHT Saranac		
RECOVERY Saranac		WINTER HARI	DINESS	Saranac			

REFERENCES

Barnes, D.K., and C.H. Hanson, An Illustrated Summary of Genetic Traits in Tetraploid and Diploid Alfalfa, ARS Technical Bul. 1370.
Barnes, D.K., et al, Standard Tests to Characterize Pest Resistance in Alfalfa Varieties. ARS-NC-19, September 1974.
Nittler, L.W., G.W. McKee, and J.L. Newcomer, Principles and Methods of Testing Alfalfa Seed for Varietal Purity. New York Agricultural Experiment Station Bul. 807.
USDA Agricultural Handbook No. 424.

COMMENTS

- 1. Observations on first flower are subjective and variable with degree of infestation of insect pests and weather damage.
- 2. Data under item 10; measurements made on approximately 350 seedlings grown in soil after 30 days as per Nittler et al, 1964.